

AD _____

MIPR NUMBER: 90MM0586

TITLE: The Effects Upon Structure of Optical Grade Polycarbonate of Various Protective Substances Applied to and Incorporated Within It

SUBTITLE: The Medical Implications of Intraocular Ballistic and Laser Protective Spectacles

PRINCIPAL INVESTIGATOR:

MAJ Brian Cavallaro, MC; MAJ Rodney Hollifield, MC;
LTC Thomas Ward, MC; COL Fleming Wertz, MC

CONTRACTING ORGANIZATION: Walter Reed Army Medical Center
Washington, DC 20307-5001

REPORT DATE: March 1995

TYPE OF REPORT: Final

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19960904 035

DTIC QUALITY INSPECTED 1


REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 1995	3. REPORT TYPE AND DATES COVERED Final (1 Jun 90 - 28 Feb 95)	
4. TITLE AND SUBTITLE The Effects Upon Structure of Optical Grade Polycarbonate of Various Protective Substances Applied to and Incorporated Within It SUBTITLE: The Medical Implications of Intraocular ...			5. FUNDING NUMBERS 90MM0586	
6. AUTHOR(S) MAJ Brian Cavallaro, MC, MAJ Rodney Hollifield, MC LTC Thomas Ward, MC, COL Fleming Wertz, MC				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Medical Center Washington, Dc 20307-5001			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER Protocol 23 ¹⁸	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 PURPOSE: We designed a study to determine the effects of intraocular fragments of ballistic protective lenses, laser protective frontserts, and the laser absorbing dyes employed in the BLPS system. BACKGROUND: The intraocular toxicity of polycarbonate, the laser absorbing dyes, and the scratch resistant coatings employed in BLPS are largely unknown. Knowledge of these potential toxicities is important in establishing medical evacuation priorities and surgical management. METHODS: Fragments of the BLPS lenses, frontserts, and suspensions of the dyes were surgically implanted into the eyes of rabbits. The animals were randomized into ten groups including positive and negative controls. The animals were examined clinically and electrophysiologically prior to implantation and enucleation. They were euthanized at various intervals up to 26 weeks. RESULTS: Those eyes implanted with 3 λ frontsert fragments exhibited supernormal dark-adapted ERG b- wave amplitudes. Intraocular inflammation was significantly greater at early time points in both the clear polycarbonate and 3 λ groups. Those eyes injected with lambda 1 and lambda 3 absorbing dyes exhibited depressed dark-adapted ERG b-wave amplitudes. CONCLUSIONS: We have developed a reliable and reproducible system for evaluating the intraocular toxicities of ballistic and laser protective eyewear. The lambda 1 and lambda 3 absorbing dyes appear to interfere with normal retinal physiology at one week. These changes are consistent with the apparent toxicity noted in the studies employing BLPS frontsert fragments in which the fragment containing all three dyes produced significant ERG abnormalities at 1 to 2 weeks. The apparent early toxicity of the BLPS 3 λ frontsert fragments, as manifested in both the clinical exam and ERG, warrant high medical evacuation priority and urgent vitrectomy in cases of accidental intraocular implantation.				
14. SUBJECT TERMS Ballistic Laser Protective Spectacles (BLPS), polycarbonate, laser absorbing dyes, toxicity			15. NUMBER OF PAGES 17	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF COLOR PAGES WHICH DO NOT REPRODUCE LEGIBLY ON BLACK AND WHITE MICROFICHE.

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

W In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Walter C. Carroll

15 AUG 86

TABLE OF CONTENTS

Title Page	1
Purpose	2
Background	2
Methods	2
Results	3
Discussion	3
Conclusions	4
Disclaimer	4
References	5
Figure 1	
Figure 2	
Figure 3	
Figure 4	
Figure 5	
Figure 6	
Figure 7	
Figure 8	

THE MEDICAL IMPLICATIONS OF INTRAOCULAR BALLISTIC AND LASER PROTECTIVE SPECTACLES (BLPS)

**Cavallaro, Brian E., Hollifield, Rodney D., Ward, Thomas P.,
Wertz, Fleming D.**

**Ocular Trauma Research Unit, Ophthalmology Svc., Walter Reed Army
Medical Center, Washington, D.C.**

PURPOSE

We designed a study to determine the effects of intraocular fragments of ballistic protective lenses and laser protective frontserts. Furthermore we evaluated the toxicity of the laser absorbing dyes employed in the BLPS system.

BACKGROUND

The intraocular toxicity of polycarbonate, the laser absorbing dyes, and the scratch resistant coatings employed in BLPS are largely unknown. Knowledge of these potential toxicities is important in establishing medical evacuation priorities and surgical management.

METHODS

Fragments (0.4 mg) of the BLPS lenses and the frontserts were surgically implanted into eyes of Albino rabbits. The animals were randomized into six groups: clear polycarbonate lens; tinted polycarbonate lens; 2 wavelength (λ) protective frontsert; 3 λ protective frontsert; iron (positive control); and sham operated eyes (negative control). The animals were evaluated by clinical exam including intraocular pressure and bright-flash ERG prior to both fragment implantation and enucleation. The animals were euthanized at 1, 2, 4, and 26 weeks. An interim clinical exam was performed at 12 weeks on the 26 week animals. The enucleated eyes were examined and sectioned for histopathological evaluation. Suspensions of 100 ug of the dyes in 0.2 ml of Balanced Salt Solution (BSS) were injected into the mid-vitreous cavity of the right eye of Albino rabbits. Thirty-two animals were randomized onto four groups: 532 nm (λ 1) absorbing dye ; 694 nm (λ 2) absorbing dye ; 1064 nm (λ 3) absorbing dye ; and BSS (vehicle). The animals were evaluated as noted above prior to intravitreal injection and at 2, 7, and 14 days thereafter. These animals were euthanized at 2 or 14 days post-injection. The eyes were enucleated, grossly examined, and sectioned for histopathological evaluation.

BLPS (Figure 1).

BLPS Fragments (Figure 2).

Fragment Surgical Technique (FIGURE 3): A) Following a partial superotemporal limbal peritomy, a 4 mm incision was created 3 mm posterior to the limbus. B) Any vitreous prolapsing from the wound was excised. C) The fragments were then placed within the vitreous cavity using intraocular foreign body forceps under direct visualization. D) The sclera was closed using 7-0

vicryl suture. After closing the conjunctiva, a 10 mg subconjunctival injection of gentamicin was administered.

RESULTS

FIGURE 4: Those eyes implanted with 3λ frontsert fragments exhibited supernormal dark-adapted ERG b-wave amplitudes when compared to sham operated eyes at 2 weeks (* $P < 0.05$). Data presented as means. Error bars are omitted for clarity. $n=5$ at each point, except at 6 months, where $n=3$.

FIGURES 5: Anterior chamber flare and vitreous cellular reaction were significantly greater at early time points (≤ 2 weeks) in those eyes implanted with either clear polycarbonate lens or 3λ frontsert fragments when compared to negative controls (* $P < 0.05$).

FIGURE 6: Fundus photographs of clear polycarbonate lens (A) and 2λ frontsert (B) fragments at two weeks, demonstrating fragment location within the vitreous cavity.

FIGURE 7: Photograph of a gross specimen containing a 2λ frontsert fragment. Enucleation occurred at two weeks. Note the flakes of frontsert coatings within the vitreous (arrow).

FIGURE 8: Photomicrograph of a Hematoxylin and Eosin stained section of an eye which had been implanted with a 3λ frontsert fragment. Enucleation occurred at two weeks. Original magnification 15X. Note the fibrous encapsulation of the frontsert fragment. F-fragment; AC-anterior chamber; V-vitreous; S-sclera.

Those eyes injected with the lambda 1 and lambda 3 absorbing dyes exhibited depressed dark-adapted ERG b-wave amplitudes at 7 days ($P=0.052$) when compared to the noninjected fellow eyes.

DISCUSSION

Supernormal ERG a- and b-wave amplitudes have been reported to occur early in toxic conditions such as metallosis bulbi (1). Components of the BLPS system which could potentially cause intraocular toxicity include polycarbonate, silica-based scratch-resistant coatings, acrylic

dye carriers, and protective dyes. Intraocular polycarbonate has been shown to be inert (2). Unwetted silicone intraocular lenses have been reported to stimulate an inflammatory encapsulating response (3). Intraocular toxicities of acrylic carriers and protective dyes are unknown. Radical formation is a potential mechanism of injury (4). Compounds related to the laser-protective dyes have been shown to generate radicals (5,6,7).

CONCLUSIONS

We have developed a reliable and reproducible system for evaluating the intraocular toxicities of ballistic and laser protective eyewear.

The lambda 1 and lambda 3 absorbing dyes appear to interfere with normal retinal physiology at one week. These changes are consistent with the apparent toxicity noted in the studies employing BLPS frontsert fragments in which the fragment containing all three dyes produced significant ERG abnormalities at 1 to 2 weeks.

The apparent early toxicity of the BLPS 3 λ frontsert fragments, as manifested in both the clinical exam and ERG, warrant high medical evacuation priority and urgent vitrectomy in cases of accidental intraocular implantation.

Further studies are needed to elucidate the pathophysiological mechanisms involved.

DISCLAIMER

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of Army or Department of Defense.

REFERENCES

1. Knave B. The ERG and Ophthalmological Changes in Experimental Metallosis in the Rabbit: II. Effects of steel, copper and aluminum particles. Acta Ophthalmologica. 1970; 48:159.
2. Yalon M, Goldberg EP, Osborn D, Stacholy J, and Sheets JW. Polycarbonate Intraocular Lenses. Journal of Cataract and Refractive Surgery. 1988; 14:393.
3. Habal MB. The Biologic Basis for the Clinical Application of the Silicones: A Correlate to Their Biocompatibility. Archives of Surgery. 1984; 119:843.
4. Southorn PA and Powis G. Free Radicals in Medicine: II. Involvement in Human Disease. Mayo Clinic Proceedings. 1988; 63:390.
5. Younes M and Strubelt, O. Vanadate-Induced Toxicity Towards Isolated Perfused Rat Livers: The Role of Lipid Peroxidation. Toxicology. 1991; 66:63.
6. Gottsch JD, Graham CR, Hairston RJ, Chen C, Green WR, and Stark WJ. Protoporphyrin IX Photosensitization of Corneal Endothelium. Archives of Ophthalmology. 1989; 107:1497.
7. Corbett MD and Corbett BR. Nucleic Acid Binding of Arylamines During the Respiratory Burst of Granulocytes. Chemical Research in Toxicology. 1988; 1:356.

This research was supported by U.S. Army Medical Research and Development Command MIPR# 90MM05863






Figure 1




1mm

Figure 2

Figure 3



Figure 5



Vitreous Cellular Reaction





Figure 6


A



B



Figure 7






Figure 8